

CROP	RATES L/ha	TIMING	WATER L/ha
Apples,	3-5	Commence application 4-6 weeks after petal fall and repeat as necessary at 10-14 day intervals	
Pears		Commence application 4-6 weeks after petal fall and repeat as necessary at 10-14 day intervals	500-1000
Apricots, Cherries, Peaches, Plums, Nectarines	3-5	Commence applications at stone hardening. Repeat applications may be made at 10-14 day intervals up to 1 week before harvest.	500-1000
Brassicas, Brussel Sprouts, Cabbage, Cauliflower, Calabrese, Broccoli	3-5	Commence application at 4-6 true leaf stage and repeat as necessary at 7-10 day intervals	
Carrots	3-5	When sufficient leaf area to intercept spray with 10-14 days between applications	250-750
Citrus	3-5	Commence application after fruit set and repeat as necessary at 14-21 day intervals	200-500 At least 1000
Cotton	2-5	3 applications at 14 day intervals commencing prior to first flower	30-100
Lucerne	5	When sufficient leaf area to intercept spray	50-250
Mango	2	Commence application at fruit swelling . N.B. Do and repeat as necessary at 14-21 day intervals not apply within 7-10 days of any oil application	400-1000 200-500
Onions	5	When sufficient leaf area to intercept spay.	400-600
Pineapple	5	Apply 4-6 weeks before harvest.	400-600
Potatoes	3-5	Apply before bulking and repeat as necessary at 7-10 day intervals.	200-500
Strawberries	5	Commence after flowering and repeat as necessary at 7-10 day intervals.	500-1000
Tomatoes	3-5	Commence application at flowering of second truss and repeat as necessary at 7-10 day intervals.	200-500
Turf	10	As soon as growth commences in spring. In the case of moderate to severe deficiency, repeat application at 14 day intervals.	500-1000
Vines	3-5	At fruit set, pea size berries, and start of ripening.	500-1000
Fertigation	Up to 50L/ha	Repeat the application at 7-14 days as necessary.	

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING



MEGA K

ANALYSIS

POTASSIUM (K) as carbonate

This product contains

Solids
Water

%W/V

g/L

41.5

415

49.00

51.00

20L 200L 1000L

Manufactured by

RUTEC PTY LTD

23 HARGRAVE STREET
TAMWORTH NSW 2340

GENERAL INSTRUCTIONS

MEGA K is a highly concentrated performance formulation to improve the yield and quality of fruit, vegetables and broadacre crops.

Potassium deficiency can occur where soil potassium levels appear sufficient, particularly during periods of peak demand, drought or water logging

MIXING AND SPRAYING

Shake the container thoroughly before use. Pour through the filter into the tank three quarters filled with water, whilst agitating. Top up with water and continue to agitate until spraying is completed.

PRECAUTIONS

Avoid application under extremes of climate eg. Rapid drying conditions, extremely slow drying conditions, frost rain or when frost or rain are anticipated

COMPATIBILITY

RUTEC products are widely compatible with many agrichemicals. Contact your distributor for information.

STORAGE AND DISPOSAL

Store in the closed original container, in a well-ventilated area as cool as possible. Protect from frost. Do not store for prolonged periods in direct sunlight.

Triple rinse containers before disposal and add hoses to tank mix or dispose of rinsate in a disposal pit away from desirable plants, their roots and watercourses.

On-site disposal of undiluted chemicals is unacceptable. Destroy empty containers by breaking, crushing or puncturing them. Dispose of the containers at a local authority landfill that does not burn its refuse. If there is no local authority landfill readily available in your area, bury the containers to a depth of 500mm or more at a licenced/approved disposal site. In some states wastes can only be buried at a licenced landfill. Do not burn empty containers or product.

SPILLAGE

Hose small spillage (less than 1 litre of concentrate) away with a large excess of water. Large spillage should not be directed to drains. Large spillage should be absorbed in sawdust or sand and the absorbent material containing the product disposed of in a sealed container at an approved disposal site.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

Do NOT contaminate streams, rivers or waterways with the chemical or used containers.

SAFETY DIRECTIONS

Strongly alkaline. Harmful if swallowed. Avoid contact with eyes, skin and clothing. Wash thoroughly with soap and water after handling. Avoid inhaling spray mist.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre Ph: 13 11 26. If in eyes, flush with water for 15 minutes and see a doctor.

EXCLUSION OF WARRANTIES AND LIMITATIONS OF LIABILITY

The purchaser acknowledges that no express or implied warranty, condition, description or representation is given or made by Rutec Pty Ltd (ABN 91 002 846 988) in relation to this product. To the maximum extent permitted by the law, all warranties, terms, conditions and any duty of care which would otherwise be implied or arise by operation of law, statute or use in relation to this product are hereby excluded as against any and all persons, whether purchasers of this product or not. To the maximum extent permitted by law, Rutec's total liability to any person in connection with this product, whether arising from a breach of any express or implied term, condition or warranty, any duty of care or otherwise, other than any liability which by these terms or otherwise is excluded, shall not exceed the amount paid by the purchaser to the vendor for this product, whether or not such liability results from or involves negli-

RUTEC